

B/S

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004665 A2(51) International Patent Classification⁷: C12P

(21) International Application Number: PCT/EP02/06566

(22) International Filing Date: 14 June 2002 (14.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

101 32 946.6 6 July 2001 (06.07.2001) DE
60/303,790 10 July 2001 (10.07.2001) US

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): DE-GUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:


— without international search report and to be republished upon receipt of that report

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): RIEPING, Mechthild [DE/DE]; Mönkebergstrasse 1, 33619 Bielefeld (DE).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY



V. 03/004665 A2

(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine.

**Process for the Preparation of L-Amino Acids using Strains
of the Enterobacteriaceae Family**

Field of the Invention

This invention relates to a process for the preparation of
5 L-amino acids, in particular L-threonine, using strains of
the Enterobacteriaceae family in which at least the *lrp*
gene is enhanced.

Prior Art

L-Amino acids, in particular L-threonine, are used in human
10 medicine and in the pharmaceuticals industry, in the
foodstuffs industry and very particularly in animal
nutrition.

It is known to prepare L-amino acids by fermentation of
strains of Enterobacteriaceae, in particular *Escherichia*
15 *coli* (*E. coli*) and *Serratia marcescens*. Because of their
great importance, work is constantly being undertaken to
improve the preparation processes. Improvements to the
process can relate to fermentation measures, such as e.g.
stirring and supply of oxygen, or the composition of the
20 nutrient media, such as e.g. the sugar concentration during
the fermentation, or the working up to the product form, by
e.g. ion exchange chromatography, or the intrinsic output
properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are
25 used to improve the output properties of these
microorganisms. Strains which are resistant to
antimetabolites, such as e.g. the threonine analogue α -
amino- β -hydroxyvaleric acid (AHV), or are auxotrophic for
metabolites of regulatory importance and produce L-amino
30 acid, such as e.g. L-threonine, are obtained in this
manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of strains of the Enterobacteriaceae family which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating the effect on the production.

Object of the Invention

The object of the invention is to provide new measures for improved fermentative preparation of L-amino acids, in particular L-threonine.

Summary of the Invention

The invention provides a process for the preparation of L-amino acids, in particular L-threonine, using microorganisms of the Enterobacteriaceae family which in particular already produce L-amino acids and in which the nucleotide sequence which codes for the lrp gene is enhanced.

Detailed Description of the Invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Threonine is particularly preferred.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or a gene or allele which codes for a corresponding enzyme or

protein with a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is
5 in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the activity or concentration of the protein in the starting microorganism.

10 The process is characterized in that the following steps are carried out:

- a) fermentation of microorganisms of the Enterobacteriaceae family in which the *lrp* gene is enhanced,
- 15 b) concentration of the corresponding L-amino acid in the medium or in the cells of the microorganisms of the Enterobacteriaceae family, and
- c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its
20 entirety or portions (> 0 to 100%) thereof optionally remaining in the product.

The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, optionally starch, optionally
25 cellulose or from glycerol and ethanol. They are representatives of the Enterobacteriaceae family chosen from the genera *Escherichia*, *Erwinia*, *Providencia* and *Serratia*. The genera *Escherichia* and *Serratia* are preferred. Of the genus *Escherichia* the species *Escherichia*
30 *coli* and of the genus *Serratia* the species *Serratia marcescens* are to be mentioned in particular.

Suitable strains, which produce L-threonine in particular, of the genus *Escherichia*, in particular of the species *Escherichia coli*, are, for example

5 *Escherichia coli* TF427
 Escherichia coli H4578
 Escherichia coli KY10935
 Escherichia coli VNIIgenetika MG442
 Escherichia coli VNIIgenetika M1
 Escherichia coli VNIIgenetika 472T23
10 *Escherichia coli* BKIIM B-3996
 Escherichia coli kat 13
 Escherichia coli KCCM-10132.

Suitable L-threonine-producing strains of the genus *Serratia*, in particular of the species *Serratia marcescens*,
15 are, for example

Serratia marcescens HNr21
 Serratia marcescens TLR156
 Serratia marcescens T2000.

Strains from the Enterobacteriaceae family which produce L-
20 threonine preferably have, inter alia, one or more genetic or phenotypic features chosen from the group consisting of: resistance to α -amino- β -hydroxyvaleric acid, resistance to thialysine, resistance to ethionine, resistance to α -methylserine, resistance to diaminosuccinic acid,
25 resistance to α -aminobutyric acid, resistance to borrelidin, resistance to rifampicin, resistance to valine analogues, such as, for example, valine hydroxamate, resistance to purine analogues, such as, for example, 6-dimethylaminopurine, a need for L-methionine, optionally a
30 partial and compensable need for L-isoleucine, a need for meso-diaminopimelic acid, auxotrophy in respect of threonine-containing dipeptides, resistance to L-threonine, resistance to L-homoserine, resistance to L-lysine, resistance to L-methionine, resistance to L-glutamic acid,

resistance to L-aspartate, resistance to L-leucine,
resistance to L-phenylalanine, resistance to L-serine,
resistance to L-cysteine, resistance to L-valine,
sensitivity to fluoropyruvate, defective threonine
5 dehydrogenase, optionally an ability for sucrose
utilization, enhancement of the threonine operon,
enhancement of homoserine dehydrogenase I-aspartate kinase
I, preferably of the feed back resistant form, enhancement
of homoserine kinase, enhancement of threonine synthase,
10 enhancement of aspartate kinase, optionally of the feed
back resistant form, enhancement of aspartate semialdehyde
dehydrogenase, enhancement of phosphoenol pyruvate
carboxylase, optionally of the feed back resistant form,
enhancement of phosphoenol pyruvate synthase, enhancement
15 of transhydrogenase, enhancement of the RhtB gene product,
enhancement of the RhtC gene product, enhancement of the
YfiK gene product, enhancement of a pyruvate carboxylase,
and attenuation of acetic acid formation.

It has been found that microorganisms of the
20 Enterobacteriaceae family produce L-amino acids, in
particular L-threonine, in an improved manner after
enhancement, in particular over-expression, of the lrp
gene.

The use of endogenous genes is in general preferred.
25 "Endogenous genes" or "endogenous nucleotide sequences" are
understood as meaning the genes or nucleotide sequences
present in the population of a species.

The nucleotide sequences of the genes of Escherichia coli
belong to the prior art and can also be found in the genome
30 sequence of Escherichia coli published by Blattner et al.
(Science 277: 1453-1462 (1997)).

The following information on the lrp gene is known, inter
alia, from the prior art:

Description: Regulator Lrp for the leucine Lrp regulon and high-affinity transport systems of branched-chain amino acids (leucine-responsive regulatory protein)

5 Reference: Willins et al.; Journal of Biological Chemistry 266(17): 10768-10774 (1991) Wang et al.; Journal of Bacteriology 176(7): 1831-1839 (1994) Friedberg et al.; Journal of Bacteriology 177(6): 1624-1626 (1995)

10 Calvo and Matthews; Microbiological Reviews 58(3): 466-490 (1994) Azam et al.; Journal of Bacteriology 181(20): 6361-6370 (1999)

Accession No.: AE000191

Alternative gene names: ihb, livR, lss, lstR, oppI, rblA,

15 mbf

The nucleic acid sequences can be found in the databanks of the National Center for Biotechnology Information (NCBI) of the National Library of Medicine (Bethesda, MD, USA), the nucleotide sequence databank of the European Molecular

20 Biologies Laboratories (EMBL, Heidelberg, Germany or Cambridge, UK) or the DNA databank of Japan (DDBJ, Mishima, Japan).

Alleles of the lrp gene which result from the degeneracy of the genetic code or due to "sense mutations" of neutral

25 function can furthermore be used.

To achieve an enhancement, for example, expression of the genes or the catalytic properties of the proteins can be increased. The two measures can optionally be combined.

To achieve an over-expression, the number of copies of the

30 corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is

additionally possible to increase the expression in the course of fermentative L-threonine production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is
5 also enhanced by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome.
Alternatively, an over-expression of the genes in question
10 can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Chang and Cohen (Journal of Bacteriology 134: 1141-1156 (1978)), in Hartley and Gregori (Gene 13:
15 347-353 (1981)), in Amann and Brosius (Gene 40: 183-190 (1985)), in de Broer et al. (Proceedings of the National Academy of Sciences of the United States of America 80: 21-25 (1983)), in LaVallie et al. (BIO/TECHNOLOGY 11: 187-193 (1993)), in PCT/US97/13359, in Llosa et al. (Plasmid
20 26: 222-224 (1991)), in Quandt and Klipp (Gene 80: 161-169 (1989)), in Hamilton (Journal of Bacteriology 171: 4617-4622 (1989)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191-195 (1998)) and in known textbooks of genetics and molecular biology.

25 Plasmid vectors which are capable of replication in Enterobacteriaceae, such as e.g. cloning vectors derived from pACYC184 (Bartolomé et al.; Gene 102: 75-78 (1991)), pTrc99A (Amann et al.; Gene 69: 301-315 (1988)) or pSC101 derivatives (Vocke and Bastia, Proceedings of the National
30 Academy of Sciences USA 80(21): 6557-6561 (1983)) can be used. A strain transformed with a plasmid vector, wherein the plasmid vector carries at least one nucleotide sequence which codes for the lrp gene, can be employed in a process according to the invention.

It is also possible to transfer mutations which affect the expression of the particular gene into various strains by sequence exchange (Hamilton et al. (Journal of Bacteriology 171: 4617 - 4622 (1989))), conjugation or transduction.

- 5 It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, with strains of the Enterobacteriaceae family to enhance one or more enzymes of the known threonine biosynthesis pathway or enzymes of anaplerotic metabolism or enzymes for the production of
10 reduced nicotinamide adenine dinucleotide phosphate, in addition to the enhancement of the *lrp* gene.

Thus, for example, one or more of the genes chosen from the group consisting of

- the *thrABC* operon which codes for aspartate kinase,
15 homoserine dehydrogenase, homoserine kinase and threonine synthase (US-A-4,278,765),
- the *pyc* gene which codes for pyruvate carboxylase (DE-A-19 831 609),
- the *pps* gene which codes for phosphoenol pyruvate
20 synthase (Molecular and General Genetics 231(2): 332-336 (1992)),
- the *ppc* gene which codes for phosphoenol pyruvate carboxylase (Gene 31: 279-283 (1984)),
- the *pntA* and *pntB* genes which code for transhydrogenase
25 (European Journal of Biochemistry 158: 647-653 (1986)),
- the *rhtB* gene which imparts homoserine resistance (EP-A-0 994 190),
- the *mgo* gene which codes for malate:quinone oxidoreductase (DE 100 348 33.5),

- the rhtC gene which imparts threonine resistance (EP-A-1 013 765),
- the thrE gene of *Corynebacterium glutamicum* which codes for the threonine export protein (DE 100 264 94.8),
- 5 • the gdhA gene which codes for glutamate dehydrogenase (Nucleic Acids Research 11: 5257-5266 (1983); Gene 23: 199-209 (1983)),
- the hns gene which codes for the DNA-binding protein HLP-II (Molecular and General Genetics 212(2): 199-202
10 (1988), Accession No. AE000222),
- the dps gene which codes for the global regulator Dps (Genes & Development 6(12B): 2646-2654 (1992), Accession No. AE000183),
- the pgm gene which codes for phosphoglucomutase (Journal
15 of Bacteriology 176: 5847-5851 (1994), Accession No. AE000172),
- the fba gene which codes for fructose bisphosphate aldolase (Biochemical Journal 257: 529-534 (1989), Accession No. AE000376),
- 20 • the ptsG gene which codes for the glucose-specific IIBC component of the phosphotransferase system PTS (Journal of Biological Chemistry 261(35): 16398-16403 (1986), Accession No. AE000210),
- the ptsH gene of the ptsHIcrr operon which codes for the
25 phosphohistidine protein hexose phosphotransferase of the phosphotransferase system PTS (Journal of Biological Chemistry 262(33): 16241-16253 (1987), Accession No. AE000329),
- the ptsI gene of the ptsHIcrr operon which codes for
30 enzyme I of the phosphotransferase system PTS (Journal

of Biological Chemistry 262(33): 16241-16253 (1987),
Accession No. AE000329),

- the crr gene of the ptsHICrr operon which codes for the glucose-specific IIA component of the phosphotransferase system PTS (Journal of Biological Chemistry 262(33):
5 16241-16253 (1987), Accession No. AE000329),
- the mopB gene which codes for chaperone GroES (Journal of Biological Chemistry 261(26): 12414-12419 (1986),
Accession No. AE000487),
- 10 • the ahpC gene of the ahpCF operon which codes for the small subunit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences USA 92(17): 7617-7621 (1995), Accession No. AE000166),
- the ahpF gene of the ahpCF operon which codes for the
15 large subunit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences USA 92(17): 7617-7621 (1995), Accession No. AE000166),

can be enhanced, in particular over-expressed.

The use of endogenous genes is in general preferred.

- 20 It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to the enhancement of the lrp gene, for one or more of the genes chosen from the group consisting of
- the tdh gene which codes for threonine dehydrogenase
25 (Journal of Bacteriology 169: 4716-4721 (1987)),
- the mdh gene which codes for malate dehydrogenase (E.C. 1.1.1.37) (Archives in Microbiology 149: 36-42 (1987)),
- the gene product of the open reading frame (orf) yjfa (Accession Number AAC77180 of the National Center for
Biotechnology Information (NCBI, Bethesda, MD, USA)),

- the gene product of the open reading frame (orf) ytfP (Accession Number AAC77179 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the pckA gene which codes for the enzyme phosphoenolpyruvate carboxykinase (Journal of Bacteriology 172: 7151-7156 (1990)),
- the poxB gene which codes for pyruvate oxidase (Nucleic Acids Research 14(13): 5449-5460 (1986)),
- the aceA gene which codes for the enzyme isocitrate lyase (Journal of Bacteriology 170: 4528-4536 (1988)),
- the dgsA gene which codes for the DgsA regulator of the phosphotransferase system (Bioscience, Biotechnology and Biochemistry 59: 256-251 (1995)) and is also known under the name of the mlc gene,
- the fruR gene which codes for the fructose repressor (Molecular and General Genetics 226: 332-336 (1991)) and is also known under the name of the cra gene, and
- the rpoS gene which codes for the sigma³⁸-factor (WO 01/05939) and is also known under the name of the katF gene,

to be attenuated, in particular eliminated or for the expression thereof to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding enzyme (protein) or gene, and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to the enhancement of the *lrp* gene, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms produced according to the invention can be cultured in the batch process (batch culture), the fed batch (feed process) or the repeated fed batch process (repetitive feed process). A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and optionally cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as

e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as
5 a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium
10 phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-
15 containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be
20 employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

25 Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can
30 be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are
35 introduced into the culture. The temperature of the

culture is usually 25°C to 45°C, and preferably 30°C to 40°C. Culturing is continued until a maximum of L-amino acids or L-threonine has formed. This target is usually reached within 10 hours to 160 hours.

- 5 The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry 30: 1190-1206 (1958)), or it can take place by reversed phase HPLC as described by Lindroth et al.
- 10 (Analytical Chemistry 51: 1167-1174 (1979)).

The process according to the invention is used for the fermentative preparation of L-amino acids, such as, for example, L-threonine, L-isoleucine, L-valine, L-methionine, L-homoserine and L-lysine, in particular L-threonine.

- 15 The present invention is explained in more detail in the following with the aid of embodiment examples.

- The minimal (M9) and complete media (LB) for Escherichia coli used are described by J.H. Miller (A Short Course in Bacterial Genetics (1992), Cold Spring Harbor Laboratory
- 20 Press). The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, ligation, Klenow and alkaline phosphatase treatment are carried out by the method of Sambrook et al. (Molecular Cloning - A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press). Unless
- 25 described otherwise, the transformation of Escherichia coli is carried out by the method of Chung et al. (Proceedings of the National Academy of Sciences of the United States of America (1989) 86: 2172-2175).

- The incubation temperature for the preparation of strains
- 30 and transformants is 37°C.

Example 1Construction of the expression plasmid pTrc99A_{lrp}

The *lrp* gene from *E. coli* K12 is amplified using the polymerase chain reaction (PCR) and synthetic
5 oligonucleotides. Starting from the nucleotide sequence of the *lrp* gene in *E. coli* K12 MG1655 (Accession Number AE000191, Blattner et al. (Science 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany). The sequences of the primers are modified such
10 that recognition sites for restriction enzymes are formed. The recognition sequence for XbaI is chosen for the *lrp*₁ primer and the recognition sequence for HindIII for the *lrp*₂ primer, which are marked by underlining in the nucleotide sequence shown below:

15 *lrp*₁: 5' - CCGGTCTAGATAAAGGAATACAGAG - 3' (SEQ ID No. 1)

*lrp*₂: 5' - CCGAAGCTTTTCCGTGTTAGCG - 3' (SEQ ID No. 2)

The chromosomal *E. coli* K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany).
20 A DNA fragment approx. 550 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA). The PCR product is
25 cleaved with the restriction enzymes XbaI and HindIII and ligated with the vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden), which has been digested with the enzymes XbaI and HindIII. The *E. coli* strain XL1-Blue MRF⁺ (Stratagene, La Jolla, USA) is transformed with the
30 ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by

control cleavage with the enzymes MluI, PvuI and SspI. The plasmid is called pTrc99Alrp (figure 1).

Example 2

Preparation of L-threonine with the strain MG442/pTrc99Alrp

- 5 The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid
10 pTrc99Alrp described in example 1 and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99Alrp and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal
15 medium with the following composition: 3.5 g/l $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 1.5 g/l KH_2PO_4 , 1 g/l NH_4Cl , 0.1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture
20 medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden,
25 Switzerland). 250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for
30 48 hours at 37°C. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer

from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99Alrp	4.7	3.6

10

Brief Description of the Figure:

- Figure 1: Map of the plasmid pTrc99Alrp containing the lrp gene.

The length data are to be understood as approx. data. The abbreviations and designations used have the following meaning:

- Amp: Ampicillin resistance gene
- lacI: Gene for the repressor protein of the trc promoter
- 20 • Ptrc: trc promoter region, IPTG-inducible
- lrp: Coding region of the lrp gene

- 5S: 5S rRNA region
- rrnBT: rRNA terminator region

The abbreviations for the restriction enzymes have the following meaning

- 5 • HindIII: Restriction endonuclease from *Haemophilus influenzae*
- MluI: Restriction endonuclease from *Micrococcus luteus* IFO 12992
- PvuI: Restriction endonuclease from *Proteus vulgaris*
- 10 • SspI: Restriction endonuclease from *Sphaerotilus* species ATCC 13925
- XbaI: Restriction endonuclease from *Xanthomonas campestris*

What is claimed is:

1. Process for the preparation of L-amino acids, in particular L-threonine, wherein the following steps are carried out:
 - 5 a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which the lrp gene or the nucleotide sequence which codes for this is enhanced, in particular over-expressed,
 - 10 b) concentration of the desired L-amino acid in the medium or in the cells of the microorganisms, and
 - c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100%) thereof
 - 15 optionally remaining in the product.
2. Process according to claim 1, wherein microorganisms in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
- 20 3. Process according to claim 1, wherein microorganisms in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partly eliminated are employed.
4. Process according to claim 1, wherein the expression of
- 25 the polynucleotide which codes for the lrp gene is increased.
5. Process according to claim 1, wherein the regulatory and/or catalytic properties of the polypeptide (protein) for which the polynucleotide lrp codes are
- 30 improved or increased.

6. Process according to claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 6.1 the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase,
 - 6.2 the pyc gene which codes for pyruvate carboxylase,
 - 6.3 the pps gene which codes for phosphoenol pyruvate synthase,
 - 6.4 the ppc gene which codes for phosphoenol pyruvate carboxylase,
 - 6.5 the pntA and pntB genes which code for transhydrogenase,
 - 6.6 the rhtB gene which imparts homoserine resistance,
 - 6.7 the mqo gene which codes for malate:quinone oxidoreductase,
 - 6.8 the rhtC gene which imparts threonine resistance,
 - 6.9 the thrE gene which codes for the threonine export protein,
 - 6.10 the gdhA gene which codes for glutamate dehydrogenase,
 - 6.11 the hns gene which codes for the DNA-binding protein HLP-II,

- 6.12 the dps gene which codes for the global regulator Dps
- 6.13 the pgm gene which codes for phosphoglucomutase,
- 5 6.14 the fba gene which codes for fructose biphosphate aldolase,
- 6.15 the ptsG gene which codes for the glucose-specific IIBC component,
- 10 6.16 the ptsH gene which codes for the phosphohistidine protein hexose phosphotransferase,
- 6.17 the ptsI gene which codes for enzyme I of the phosphotransferase system,
- 15 6.18 the crr gene which codes for the glucose-specific IIA component,
- 6.19 the mopB gene which codes for chaperone GroES,
- 6.20 the ahpC gene which codes for the small sub-unit of alkyl hydroperoxide reductase,
- 20 6.21 the ahpF gene which codes for the large sub-unit of alkyl hydroperoxide reductase,

is or are enhanced, in particular over-expressed, are fermented.


- 7. Process according to claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:

7.1 the adh gene which codes for the alcohol dehydrogenase,

- 7.2 the mdh gene which codes for malate dehydrogenase,
- 7.3 the gene product of the open reading frame (orf) yjfa,
- 5 7.4 the gene product of the open reading frame (orf) ytfP,
- 7.5 the pckA gene which codes for phosphoenol pyruvate carboxykinase,
- 7.6 the poxB gene which codes for pyruvate oxidase,
- 10 7.7 the aceA gene which codes for isocitrate lyase,
- 7.8 the dgsA gene which codes for the DgsA regulator of the phosphotransferase system,
- 7.9 the fruR gene which codes for the fructose repressor,
- 15 7.10 the rpoS gene which codes for the sigma³⁸ factor

is or are attenuated, in particular eliminated or reduced in expression, are fermented.

Figure 1:



SEQUENCE PROTOCOL

5 <110> Degussa AG
<120> Process for the preparation of L-amino acids using strains of the Enterobacteriaceae family
10 <130> 020093 BT
<160> 2
<170> PatentIn version 3.1
15 <210> 1
<211> 24
<212> DNA
<213> artificial sequence
20 <220>
<221> Primer
<222> (1)..(24)
<223> lrp1
25 <400> 1
ccggtctaga taaggaatac agag 24
<210> 2
<211> 22
30 <212> DNA
<213> artificial sequence
<220>
<221> Primer
35 <222> (1)..(22)
<223> lrp2
<400> 2
ccgaagcttt tccgtgtag cg 22
40 22

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004665 A3

(51) International Patent Classification⁷: C12N 15/31,
C12P 13/08

(21) International Application Number: PCT/EP02/06566

(22) International Filing Date: 14 June 2002 (14.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
101 32 946.6 6 July 2001 (06.07.2001) DE
60/303,790 10 July 2001 (10.07.2001) US

(71) Applicant (*for all designated States except US*): DE-
GUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düssel-
dorf (DE).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): RIEPING, Mechthild
[DE/DE]; Mönkebergstrasse 1, 33619 Bielefeld (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR). OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) *for US only*


Published:

— with international search report

(88) Date of publication of the international search report:
21 August 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY



(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine.

WO 03/004665 A3

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 02/06566

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/31 C12P13/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 090 993 A (DEGUSSA) 11 April 2001 (2001-04-11) claims 7,8	1-7
A	US 4 278 765 A (DEBABOV VLADIMIR G ET AL) 14 July 1981 (1981-07-14) cited in the application the whole document	1-7
A	ERNSTING B R ET AL: "CHARACTERIZATION OF THE REGULON CONTROLLED BY THE LEUCINE-RESPONSIVE REGULATORY PROTEIN IN ESCHERICHIA-COLI" JOURNAL OF BACTERIOLOGY, vol. 174, no. 4, 1992, pages 1109-1118, XP001150347 ISSN: 0021-9193 the whole document	1-7

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *G* document member of the same patent family

Date of the actual completion of the international search

16 April 2003

Date of mailing of the international search report

28/04/2003

Name and mailing address of the ISA

European Patent Office, NL 5016 FR Utrecht, NL
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lejeune, R

INTERNATIONAL SEARCH REPORT

International Publication No

PCT/EP 02/06566

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LAURENT-WINTER CHRISTINE ET AL: "Role of Escherichia coli histone-like nucleotide-structuring protein in bacterial metabolism and stress response: Identification of targets by two-dimensional electrophoresis." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 244, no. 3, 1997, pages 767-773, XP001150348 ISSN: 0014-2956 the whole document	1-7
A	EP 0 994 190 A (AJINOMOTO KK) 19 April 2000 (2000-04-19) the whole document	1-7
A	SAWERS G: "THE ANAEROBIC DEGRADATION OF L-SERINE AND L-THREONINE IN ENTEROBACTERIA: NETWORKS AND PATHWAYS AND REGULATORY SIGNALS" ARCHIVES OF MICROBIOLOGY, BERLIN, DE, vol. 171, no. 1, 1998, pages 1-5, XP002953871 ISSN: 0302-8933 the whole document	1-7
E	WO 03 004598 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 16 January 2003 (2003-01-16) claim 6 & WO 03 004663 A 16 January 2003 (2003-01-16) & WO 03 004664 A & WO 03 004669 A & WO 03 004670 A & WO 03 004671 A & WO 03 004674 A & WO 03 004675 A	6
T	WO 03 004662 A (DEGUSSA) 16 January 2003 (2003-01-16) the whole document	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Publication No

PCT/EP 02/06566

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1090993	A	11-04-2001	DE 19947792 A1	12-04-2001
			AU 6245100 A	12-04-2001
			BR 0004642 A	12-06-2001
			CA 2319722 A1	05-04-2001
			CN 1290746 A	11-04-2001
			EP 1090993 A1	11-04-2001
			HU 0003894 A2	28-10-2002
			JP 2001128692 A	15-05-2001
			PL 342985 A1	09-04-2001
			SK 14592000 A3	11-06-2001
			US 2003017555 A1	23-01-2003
US 4278765	A	14-07-1981	SU 875663 A1	15-09-1982
			HU 190999 B	28-12-1986
EP 0994190	A	19-04-2000	RU 2144564 C1	20-01-2000
			AU 4755099 A	20-04-2000
			BR 9904955 A	12-12-2000
			CN 1254014 A	24-05-2000
			EP 0994190 A2	19-04-2000
			JP 2000116390 A	25-04-2000
			KR 2000029006 A	25-05-2000
			SK 140899 A3	16-05-2000
			US 6303348 B1	16-10-2001
			US 2002102670 A1	01-08-2002
			US 2002058314 A1	16-05-2002
			ZA 9906042 A	04-04-2000
WO 03004598	A	16-01-2003	DE 10132946 A1	16-01-2003
			WO 03004663 A2	16-01-2003
			WO 03004669 A2	16-01-2003
			WO 03004674 A2	16-01-2003
			WO 03004670 A2	16-01-2003
			WO 03004664 A2	16-01-2003
			WO 03004598 A2	16-01-2003
			WO 03004665 A2	16-01-2003
			WO 03004671 A2	16-01-2003
			WO 03004675 A2	16-01-2003
WO 03004662	A	16-01-2003	DE 10132945 A1	16-01-2003
			WO 03004662 A2	16-01-2003
			US 2003017556 A1	23-01-2003